

CHAPTER ONE

OUR OWN FACE VERSUS  
OUR PARTNER'S FACE:  
NEURAL DIFFERENTIATIONS

MARIE ARSALIDOU,<sup>1</sup>  
EMMANUEL J. BARBEAU,<sup>2</sup> SARAH J BAYLESS<sup>3</sup>  
AND MARGOT J. TAYLOR<sup>1</sup>

<sup>1</sup>UNIVERSITY OF TORONTO, CANADA;

<sup>2</sup>CNRS, TOULOUSE, FRANCE;

<sup>3</sup>WINCHESTER UNIVERSITY, ENGLAND

Despite the fact that faces share a basic configuration, we are very skilled at recognizing hundreds of different faces. Once faces are known, they are linked with semantic and social information which together become the means for distinguishing a person. This is referred to as “person knowledge” which contains information such as social relations and personal traits (Gobbini and Haxby 2007).

The ability to readily process information about a person we encounter influences how we interact with him or her. The human brain processes faces of conspecifics with great efficiency (Dufour, Pascalis and Petit 2006) and functional imaging has been a useful tool for identifying the location of brain functions such that quantitative measurements of activity can be related to the brain structure. There is general agreement among neuroscientists that photographs of faces recruit specific brain areas, including the fusiform gyri, located on the inferior surface of the temporal lobes (e.g., Puce et al. 1996; McCarthy et al. 1997; Fusar-Poli et al. 2009 for a meta-analysis). Although the fusiform gyri are key for the visual recognition of faces (e.g., McCarthy et al. 1997; Eger et al. 2005), the semantic recognition of faces is not limited to their visual characteristics. Familiar faces carry a host of non-visual information, information based on previous social interactions, which renders them higher in social and

emotional salience to the viewer. Therefore, current face processing models accept that faces engage a network of brain areas (Ishaii, Haxby and Ungerleider 2002; Ishaii, Schmidt and Boesiger 2005; Gobbini and Haxby 2007; Taylor, Arsalidou, Bayless, Morris, Evans and Barbeau 2009).

Familiarity can change the neural response to faces beyond the simple visual memory of the appearance of a face (Gobbini et al. 2004). Although, there is a general agreement that familiarity modulates activation in the brain (e.g., Arsalidou, Barbeau, Bayless, Taylor 2010; Leveroni et al. 2000; Taylor et al. 2009), a methodological shortcoming in this line of research is the range of familiarity of the face stimuli employed. Researchers' classification of familiar face stimuli has varied from personally familiar to recently acquired familiarization. For example, photographs of friends and family (Gobbini et al. 2004) would carry long-term social interactions that range in quality; on the other hand photographs of fraternity brothers (Platek et al. 2006) would likely be recent and associated with more variable information, while faces learned in a lab setting to induce familiarity (Gobbini and Haxby 2006; Dubois et al. 1999) would carry limited or no person knowledge to the viewer. Visual familiarity does not entail personal familiarity and the extent and quality of social interactions carry different emotional significance. In addition, personally familiar faces are a special class of faces that have the advantage of being processed frequently and repeatedly, an aspect likely reflected in neural processes even at the visual level.

Neuroimaging studies have modelled the intricate neural networks involved in face recognition. Haxby, Hoffman and Gobbini (2000) initially proposed a model of face recognition that implicates two major neural systems: the core visual processing system, which includes the ventral visual pathways and fusiform gyri, and more extended systems which include limbic structures and frontal areas for processing and emotional and cognitive information contained in faces. This model was updated to include a specific focus on the components that influence recognition of familiar faces (Gobbini and Haxby 2007). The authors propose that recognition of familiar faces first recruits a spatially distributed network in the brain that not only includes areas of visual processing but areas associated with cognitive and social functions. Second, they claim that areas associated with "theory of mind" are also recruited for the spontaneous retrieval of "person knowledge" in the process of face recognition. Lastly, they suggest that social interactions and familiarity recognition modulate the emotional response to familiar faces. These three aspects (core system, person knowledge and emotion) all play a role in the

successful identification of a face. According to Gobbini and Haxby (2007), the core system in the brain includes the posterior superior temporal sulci (STS) and inferior occipital and fusiform gyri, which together assess the visual appearance of a face; the person knowledge engages the anterior paracingulate, the posterior STS, the temporo-parietal junction, anterior temporal cortex, the precuneus and the posterior cingulate; emotional processing of faces includes areas such as the amygdalae, the insula and the striatum.

Two of the most important faces in our lives are our own face and our partner's face. It is a common experience that we feel strongly about our face as well as our partner's face; however, the reasons may be very different. As the narcissus myth exemplifies, our own face is very special given that it represents our own emotional state and especially how we view ourselves in the world (i.e. self awareness). Kircher and colleagues (2001) suggest that one's own face entails self-referential processing which requires emotional and associative cognitive processes. They found that one's own face activated right limbic areas, right insula, left superior temporal gyrus, left inferior parietal gyrus and left prefrontal cortex (BA 8/9 and BA 45/46). A later study by Sugriva and colleagues (2005) also investigated the neural correlates of visual self-recognition and found that compared to an unfamiliar face, activation related to one's own face was present in the right occipito-temporo-parietal junction, right frontal operculum, left fusiform gyrus, bilateral cingulate cortex and parahippocampal gyri. Platek and colleagues (2006) reported activation in the right postcentral, supramarginal and superior temporal gyri linked to one's own face. These researchers also compared the subject's own face with another familiar face (fraternity brother), and the subject's own face activated the right superior frontal gyrus, inferior parietal and medial frontal lobes and left middle temporal gyrus. One's own face compared to a colleague's face showed activation in the right inferior frontal gyrus and the right insula (Devue et al. 2007). Digital morphs between their own face and a gender-matched familiar face elicited more activity in the right inferior parietal lobule, inferior frontal gyrus and inferior occipital gyrus for images containing more of one's face than the familiar face suggesting that this network is engaged in maintaining self/other distinctions (Uddin et al. 2005). Overall, key regions that repeatedly activate to one's own face are the left fusiform gyrus, bilateral middle and inferior frontal gyri and the right precuneus (see Platek, Wathne, Tierney & Thomson 2008 for a meta-analysis).

In contrast to the visual, cognitive and emotional information that are activated concurrently when we see our image, our partner's face would

likely activate a set of areas associated with our previous social interactions. In social neuropsychology, researchers have used the partner's face as an effective stimulus for eliciting an emotional state (Fisher, Aron & Brown 2005). This research is usually referred to as research on romantic love that investigates the neural correlates of mate choice (Fisher, Aron & Brown 2005, 2006; Bartels & Zeki 2000, 2004). Fisher and colleagues (2005) studied 17 intensely "in love" young adults who were in a relationship between 0.08 to 1.41 years (1-17 months), with a mean relationship length of 0.61 years. The main brain regions showing increased activation for pictures of their beloved were the right ventral tegmental area and right posterior-dorsal body of the caudate nucleus. Some of the participants also showed activity in the right insula, and right anterior and posterior cingulate cortex. The latter findings are in agreement with the study of Bartels & Zeki (2000), who also tested 17 young adults who were in a relationship between 0.7 to 4.1 years, with a mean relationship of 2.4 years. These intensely "in love" participants viewed photographs of their partner and generated activity bilaterally in the medial insula, the anterior cingulate, the caudate nucleus and the putamen. Bartel & Zeki (2000) also reported deactivation in the cingulate gyrus and the amygdalae as well as prefrontal, parietal and middle temporal cortices. Overall, these two studies suggest that early stages of romantic love activate areas in the limbic system when viewing a partner's face (Fisher et al. 2005), which are represented by the emotional system in the model by Gobbini & Haxby (2007). At a later stage in a relationship (Bartel & Zeki 2000) this system seems to evolve to include areas associated with person knowledge. Individuals that sustain relationships that survive many years (e.g., married ten to twenty-nine years) may also elicit activity in areas similar to new love (Acevedo, Aron, Fisher & Brown 2011).

## 1. Current Study

The current study examined the neural differentiation between one's own face and one's partner's face past the initial intense "in love" period. Familiarity accumulates naturally over years of social interactions and long exposure, thus we defined a long-term relationship as one where a couple has cohabitated for a minimum of two years ( $X = 9$  yrs). The current report was part of a larger investigation including parent and famous faces (Taylor et al. 2009) which are not discussed here.

Compared to baseline (i.e. fixation cross), we anticipated that all faces (own, partner and unfamiliar) would activate areas in the core system such

as the posterior superior temporal sulcus, the inferior occipital lobes and fusiform gyri. Based on previous studies we also expected that one's own face would generate activation in areas associated with person knowledge (i.e. self-awareness) to a greater extent than emotional processing and that this activation would be primarily in the right hemisphere. A partner's face was expected to activate areas involved in both the emotional processing system and person knowledge such as the amygdala and anterior cingulate.

## **1.1 Methods**

### **1.1.1 - Participants**

We studied ten heterosexual participants, four males, with a mean age of 35.4 years (7.7 SD), who had lived with their partner for at least two years. Average time together was  $9.1 \pm 5.01$  years.

### **1.1.2 - Stimuli**

Each subject provided his or her own digitised photographs of their partner's, their own and parents' face following a standardised protocol controlling for neutrality (no emotion on the face), gaze direction (looking straight at the camera) and light conditions. Two unfamiliar photographs and two famous photographs of faces (male and female) were also included, which were different for each participant (see Taylor et al. 2009, for greater detail). The set was processed to remove all information unrelated to the faces themselves (background, paraphernalia, etc.). The photographs were converted to greyscale and contrast level among photographs controlled. A "ghost" image (~ 20 trials) was prepared for each set superposing all photographs. Stimuli were repeated such that at least forty trials of each face type (e.g. partner, own and unfamiliar) were presented. The faces were presented in a pseudo-random order for 500 ms and inter-stimulus interval (ISI) was jittered between 1700 and 2000ms.

### **1.1.3 - Task Instructions**

Participants were asked to focus on the photographs and pressed a button for the rare "ghost" image.

### 1.1.4 - fMRI Acquisition

All MR imaging was conducted on a 1.5T Signa Twin EXCITE3 scanner (GE Medical Systems, WI; software rev.12M4) with a standard quadrature head coil. Foam padding comfortably restricted head motion. A set of high resolution T1-weighted 3D SPGR images covering the whole brain was acquired (116 slices, voxel size =  $1 \times 1 \times 1.5\text{mm}^3$ , 2 NEX, 7min) prior to the acquisition of functional images as an anatomical reference. Functional images were acquired with a standard gradient-recalled echo-planar imaging sequence (TE/TR/ $\alpha$  = 40ms/2000ms/90°, voxel size =  $3.75 \times 3.75 \times 5\text{mm}^3$ ) over 27 contiguous axial slices with interleaved acquisition.


Face stimuli were displayed on an MR compatible goggle system (CinemaVision, Resonance Technology Inc., CA). Subjects responded to trials using an MR compatible keypad (Lumitouch, Photonics Control, CA). Stimuli were controlled and responses recorded using the software *Presentation* (Neurobehavioral Systems Inc., CA) on a personal computer.

### 1.1.5 - Data Processing and Analyses

Data analyses were carried out in AFNI (Cox 1996), using motion correction, 8mm spatial smoothing (RMSD), signal intensity normalization for percent signal change. A general linear model was applied to deconvolve the data using a fixed hemodynamic response function for all participants across all face types. Images were transformed into Talairach stereotaxic space and resampled to  $3\text{mm}^3$ . Group images were analyzed using a random effects analysis of variance. The results were thresholded at  $p < 0.01$  (corrected) for the whole brain. To control for multiple comparisons we performed 1000 Monte Carlo iterations at an uncorrected  $p$ -value of 0.01 for individual voxels on 43,349 voxels, yielding a minimum volume of  $5670\mu\text{l}$  (210 voxels) at a  $p$ -value of 0.05.

## 1.2 Results

The results revealed overall differences between the two types of familiar faces (Fig. 1-1 below). Compared to baseline, partner faces activated bilateral fusiform gyri, bilateral lingual gyri, right cuneus, the right parahippocampal gyrus, right middle temporal gyrus, right superior temporal gyrus, left precuneus and left middle frontal gyrus. Pictures of the participant's face (i.e. own faces) activated bilateral fusiform gyri, bilateral inferior occipital and bilateral lingual gyrus, bilateral cuneus and right precuneus, compared to baseline.



**Fig. 1-1.** Activity related to own and partner faces when compared to (a) fixation cross and (b) unfamiliar faces.

Own and partner faces were contrasted with unfamiliar faces; the anterior cingulate, cingulate gyrus and medial frontal gyrus bilaterally and left middle frontal gyrus activated more to own faces than unfamiliar faces. There was no significant activation in the comparison between own face and partner's face. However, the partner's face showed the most extended activations compared to unfamiliar faces with bilateral activation in the anterior cingulate, cingulate and medial frontal gyri, as well as activation in the left middle frontal gyrus, left inferior frontal gyrus (BA 47), left middle temporal gyrus, left parahippocampal gyrus, left amygdala, left insula, left thalamus and left precuneus.

### 1.3 Discussion

We examined the neural differentiations of two types of personally familiar faces. One's face and one's partner's face play an important role

in our daily interactions, yet the neural correlates for processing these faces has been rarely investigated together. Several studies examined the neural responses to one's own face and a meta-analysis suggest that brain activity of self-face processing engages the fusiform gyri, as well as extensive regions associated with person knowledge (Platek et al. 2008; Gobbini & Haxby 2007). fMRI studies of the neural correlates of partner's faces cover relationships lengths of less than one year (Fisher et al. 2005), four years (Bartels and Zeki 2000), and to a maximum of twenty-one years (Acevedo et al. 2011). We recruited participants who cohabitated with their partner longer than 2 years, with a mean cohabitation length of 9.1 years. Activation related to own faces and partner's faces is discussed in relation to baseline and unfamiliar faces.

### **1.3.1 Personally Familiar Faces vs. Baseline**

Activations of personally familiar faces compared to baseline were expected to recruit areas associated with the visual recognition of faces (e.g. Puce et al. 1996; McCarthy et al. 1999; Platek et al. 2006). Bilateral activation in the fusiform gyri, as well as lingual gyri and cuneus were present for processing both one's own face and partner's face. The inferior occipital gyri were activated only for own faces. The location of these areas in the occipital lobes is suggestive of their role in visual processing (Kolb & Whishaw 2003). The precuneus, in the parietal lobes, was also activated for both one's own face and partner's face but in the opposite hemispheres, right and left respectively. Among other cognitive functions, the precuneus is associated with the visuospatial analysis of objects (Faillenot, Decety & Jeannerod 1999), a recognition network supporting retrieval (Reber, Wond & Buxton 2002; Nagahama et al. 1999) as well as person knowledge processing or face recognition (Gobbini & Haxby 2006, 2007; Bayle & Taylor 2010).

Unlike own faces, partner's faces also activated the middle and superior temporal gyri, the parahippocampal and the middle frontal gyri (BA 9). Activity in the middle frontal gyrus has been associated with cognitive functions such as working memory and attention (e.g., Petrides 1996; Christoff & Gabrieli 2000). Activation in this area was also associated with personally familiar faces (Bartel & Zeki 2000; Gobbini & Haxby 2006; Platek et al. 2006). In the present study, the middle frontal gyri were probably involved in the system responsible for holding the person information in mind. Middle temporal cortex supports recognition of familiar faces, along with the parahippocampal gyri (Barbeau et al. 2008; Leveroni et al. 2000). The middle temporal cortices, which also

activate to mother's faces (Arsalidou et al. 2010), may be related to self-related processing of personal experiences associated with the person, in this case the partner's face. The parahippocampal areas are involved in encoding visual information (e.g. Rombouts et al. 1999; Kirchoff et al. 2000), and in face recognition studies they are associated with memory retrieval of personal and social knowledge (Sugiura et al. 2005). STS areas are linked with the social cognition of faces (Allison et al. 2000) and together with the precuneus, these areas respond to personally familiar faces (Gobbini et al. 2004; Uddin et al. 2005).

Overall, compared to baseline, personally familiar faces exhibited activation primarily associated with the core system of processing faces. Partner faces activated more regions associated with the extended face processing system, which are likely associated with the mnemonic, semantic and emotional attributes of the partner.

### **1.3.2 Personally Familiar Faces vs. Unfamiliar Faces**

Comparisons with unfamiliar faces showed that core system areas including the fusiform gyri did not survive correction, as these regions respond to faces regardless of familiarity (Puce et al. 1996; McCarthy et al. 1997; Gobbini & Haxby 2007), emotional valence or mode of presentation (Arsalidou, Morris & Taylor 2011). These contrasts elicited activity in areas associated with extended affective and cognitive system, which included bilateral anterior cingulate (BA 32), cingulate gyrus, medial frontal gyri and left middle frontal activation for both own face and partner's face. The cingulate gyrus is recruited in a variety of paradigms of higher cognitive function (Petit et al. 1998), such as executive attention and set shifting (Catafau et al. 1998). Turak and colleagues (2002) concluded that the cingulate gyrus is a multimodal area involved in cognitive activity in general, including executive attention. Specifically, the anterior and dorsal parts of the cingulate gyri were found to be implicated in emotional and cognitive processes, respectively (Bush, Luu & Posner 2000 for review). In relation to face-recognition studies, the anterior cingulate gyrus has been associated with familiar faces (Platek et al. 2006; Sugiura et al. 2005) and may play a role in the integration of information (Deveaux et al. 2007). Similarly, prefrontal regions, such as the middle frontal and medial frontal gyri have been linked to executive functions such as manipulation and monitoring information held in mind (Christoff & Gabrieli 2000). Together, the anterior cingulate gyrus and prefrontal regions may facilitate the integration and monitoring or

reminiscing on personal information present in both one's own face and partner's face but not in the unfamiliar faces.

Areas responding only to partner's faces were the left inferior frontal gyrus (BA 47), insula, amygdala and thalamus. A more general function attributed to the inferior frontal gyrus is that of maintenance of a few items in mind (Christoff & Gabrieli 2000). Even though participants in this study were not explicitly asked to maintain information, the presentation of their partner's face may have implicated different memories and emotional components elicited by the pictures. The inferior frontal gyrus was also found active to mother's faces when compared to unfamiliar faces, albeit in the opposite hemisphere (Arsalidou et al. 2010). Thus, the inferior frontal regions may be part of the extended network for processing socio-cognitive aspects of a face. Emotional valence has been associated with the insula, the amygdalae and the thalamus (e.g. Britton et al. 2006). In social neuroimaging, insula activity is associated with partner's faces (Fisher et al. 2005; Bartels & Zeki 2000), as well as modulating emotions (Heinzel et al. 2005). Similarly, the amygdalae have been linked to processing of emotive states (Arsalidou et al. 2011) and social and emotional attachment to faces (Gobbini et al. 2004). Together, the amygdalae and the insula would be linked with the processing of emotions as part of face recognition (Gobbini & Haxby 2007). Compared to unfamiliar faces, own faces recruit areas primarily associated with person knowledge (i.e., self-assessment), while partner's faces recruit areas associated both with person knowledge and emotional processing.

In conclusion, humans have a complex social organization and faces play a major role in the processing of the relations an individual may have. By capitalizing on the neural correlates of own and partner faces, we showed that in long-term relationships the brain recruits areas referred to as the emotional system as well as person knowledge. Own face processing only recruited areas previously associated with person knowledge, while partner face processing recruited areas associated both with person knowledge and emotions. In a direct comparison, the neural correlates of own faces and partner faces were not significantly different. This can be attributed to the saliency of both faces in terms of person knowledge and to the small sample size, which may not be optimal in investigating the subtle differences that appeared when analyzing each face type separately. These findings contribute to the literature of face processing and romantic love, extending evidence of brain areas implicated in partner's face processing in a long-term relationship. Future experiments still need to determine the fine-grained differences between

processing personally familiar faces and the neural changes occurring across long-term exposure and familiarity.

## References

- Acevedo, B.P., A. Aron, H. E. Fisher & L. L. Brown. "Neural correlates of long-term intense romantic love. Social Cognitive and Affective Neuroscience": doi:10.1092/scan/nsq092, 2011.
- Allison, T., A. Puce & G. McCarthy. "Social Perception from Visual Cues: Role of the STS Region." *Trends in Cognitive Sciences* 4 (2000): 267–278.
- Arsalidou, M., E. J. Barbeau, S. J. Bayless & M. J. Taylor. "Brain Responses Differ to Faces of Mothers and Fathers." *Brain and Cognition* 74 (1) (2010): 47–51.
- Arsalidou, M., D. Morris & M. J. Taylor. "Converging evidence for the advantage of dynamic facial expressions." *Brain Topography* 24 (2011): 149–163.
- Barbeau, E. J., M. J. Taylor, J. Regis, P. Marquis, P. Chauvel & C. Liégeois-Chauvel. "Spatio-Temporal Dynamics of Face Recognition." *Cerebral Cortex* 18 (2008): 997–1009.
- Bartels, A. & S. Zeki. "The Neural Basis of Romantic Love." *NeuroReport* 11 (2000): 3829–3834.
- Bartels, A. & S. Zeki. "The Neural Correlated of Maternal and Romantic Love." *NeuroImage* 21 (2004): 1155–1166.
- Bayle, D. J. & M. J. Taylor. "Attention Inhibition of Early Cortical Activation to Fearful Faces." *Brain Research* 1313 (2010): 113–123.
- Britton, J. C., K. Luan Phan, S. F. Taylor, R. C. Welsh, K. C. Berridge & I. Leberzon. "Neural Correlates of Social and Nonsocial Emotions: An fMRI Study." *NeuroImage* 31 (2006): 397–409.
- Bush, G., P. Luu & M. I. "Posner. Cognitive and Emotional Influences in Anterior Cingulate Cortex." *Trends in Cognitive Sciences* 4 (2000): 215–222.
- Catafau, A. M., E. Parellada, F. Lomena, M. Bernardo, J. Setoain, S. Catarineu, J. Pavia & R. Herranz. "Role of the Cingulate Gyrus During the Wisconsin Card Sorting Test: A Single Photon Emission Computed Tomography Study in Normal Volunteers." *Psychiatric Research: Neuroimaging* 83 (1998): 67–74.
- Christoff, K. & J. D. E. Gabrieli. "The Frontopolar Cortex and Human Cognition: Evidence for a Rostocaudal Hierarchical Organization Within the Human Cortex." *Psychobiology* 28 (2000): 168–186.

- Cox, R. W. "Afni: Software for Analysis and Visualization of Functional Magnetic Resonance Neuroimages." *Computers and Biomedical Research* 29 (1996): 162–173.
- Devue, C., F. Collette, E. Balteau, C. Degueldre, C. Luxen, P. Maquet & S. Brédart. "Here I Am: The Cortical Correlates of Visual Self-Recognition." *Brain Research* 1143 (2007): 169–182.
- Dubois, S., B. Rossion, C. Schiltz, J. M. Bodart, C. Michel, R. Bruyer & M. Crommelinck. Effect of Familiarity on the Processing of Human Faces. *NeuroImage* 9 (1999): 278–289.
- Dufour, V., O. Pascalis & O. Petit. "Face Processing Limitation to Own Species in Primates: A Comparative Study in Brown Capuchins, Tonkean Macaques and Humans." *Behavioural Processes* 73 (2006): 107–113.
- Eger, E., S. R. Schweinberger, R. J. Dolan & R. N. Henson. "Familiarity Enhances Invariance of Face Representations in Human Ventral Visual Cortex: fMRI Evidence." *NeuroImage* 26 (2005): 1128–1139.
- Faillenot, I., J. Decety & M. Jeannerod. "Human Brain Activity Related to the Perception of Spatial Features of Objects." *NeuroImage* 10 (1999): 114–124.
- Fisher, H., A. Aron & L. L. Brown. "Romantic Love: An fMRI Study of a Neural Mechanism for Mate Choice." *The Journal of Comparative Neurology* 493 (2005): 58–62.
- Fisher, H., A. Aron & L. L. Brown. "Romantic Love: A Mammalian Brain System for Mate Choice." *Philosophical Transactions of the Royal Society B* 361 (2006): 2173–2186.
- Fusar-Poli, P. Placentino, A. Carletti, F. Landi, P. Allen, P. Surguladze, S. Benedetti, F. Abbamonte, M. Gasparotti, R. Barale, F. Perez, J. McGuire, P. & P. Politi. "Functional Atlas of Emotional Faces Processing: A Voxel-Based Meta-Analysis of 105 Functional Magnetic Resonance Imaging Studies." *Journal of Psychiatric Neuroscience* 34 (6) (2009): 418–432.
- Gobbini, I. M. & J. V. Haxby. "Neural Response to the Visual Familiarity of Faces." *Brain Research Bulletin* 71 (2006): 76–82.
- Gobbini, I. M. & J. V. Haxby. "Neural Systems for Recognition of Familiar Faces." *Neuropsychologia* 45 (2007): 32–41.
- Gobbini, I. M., E. Leibenluft, N. Santiago & J. V. Haxby. "Social and Emotional Attachment in the Neural Representation of Faces." *NeuroImage* 22 (2004): 1628–1635.
- Haxby, J. V., E. A. Hoffman & I. M. Gobbini. "The Distributed Human Neural System for Face Perception." *Trends in Cognitive Science* 4 (2000): 223–233.

- Heinzel, A., F. BERPPOHL, F. NIESE, R. PFENNIG, A. PASCUAL-LEONE, A. SCHLAUG, G. & G. NORTHOFF. "How do we Modulate our Emotions? Parametric fMRI Reveals Cortical Midline Structures as Regions Specifically Involved in the Processing of Emotional Valences." *Cognitive Brain Research* 25 (2005): 348–358.
- Ishai, A., J. V. Haxby & L. G. Ungerleider. "Visual Imagery of Famous Faces: Effects of Memory and Attention Revealed by fMRI." *NeuroImage* 17 (2002): 1729–1741.
- Ishai, A. C. F. Schmidt & P. Boesiger. "Face Perception is Mediated by a Distributed Cortical Network." *Brain Research Bulletin* 67 (2005): 87–93.
- Kircher, T. T. J., C. Senior, M. L. Phillips, S. Rabe-Hesketh, P. J. Benson, E. Bullmore, M. Brammer, A. Simmons, M. Bartels & A. S. David. "Recognizing One's Own Face." *Cognition* 78 (2001): B1–B15.
- Kirchhoff, B. A., A. D. Wagner, A. Maril & C. E. Stern. "Prefrontal-Temporal Circuitry for Episodic Encoding and Subsequent Memory." *The Journal of Neuroscience* 20 (2000): 1673–6180.
- Kolb, B. & I. Q. Whishaw. *Fundamentals of Human Neuropsychology* (5th ed.). New York: Worth, 2003.
- Leveroni, C. L., M. Seidenberg, A. R. Mayer, L. A. Mead, J. R. Binder & S. M. Rao. "Neural Systems Underlying the Recognition of Familiar and Newly Learned Faces." *The Journal of Neuroscience* 20 (2000): 878–886.
- McCarthy, G., A. Puce, J. C. Gore & T. Allison. "Face-Specific Processing in the Human Fusiform Gyrus." *The Journal of Cognitive Neuroscience* 9 (1997): 605–610.
- McCarthy, G., A. Puce, J. C. Gore & T. Allison. 1999. "Electrophysiological Studies of Human Face Perception. II: Response Properties of Face-specific Potentials Generated in Occipitotemporal Cortex." *Cerebral Cortex* 9 (5) (1999): 431–444.
- Nagahama, Y., T. Okada, Y. Katsumi, T. Hayashi, H. Yamauchi, N. Sawamoto, K. Toma, K. Nakamura, T. Hanakawa, J. Konishi, H. Fukuyama & H. Shibasaki. "Transient Neural Activity in the Medial Superior Frontal Gyrus and Precuneus Time Locked with Attention Shift Between Object Features." *NeuroImage* 10 (1999): 193–199.
- Petit, L., S. M. Courtney, L. G. Ungerleider & J. V. Haxby. "Sustained Activity in the Medial Wall During Working Memory Delays." *The Journal of Neuroscience* 18 (1998): 9429–9437.
- Petrides, M. "Lateral Frontal Cortical Contribution to Memory." *The Neurosciences* 8 (1996): 57–63.

- Platek, S. M., J. W. Loughhead, R. C. Gur, S. Busch, K. Ruparel, N. Phend, I. S. Panyavin & D. D. Langleben. "Neural Substrates for Functionally Discriminating Self-Face from Personally Familiar Faces." *Human Brain Mapping* 27 (2006): 91–98.
- Platek, S. M., J. Wathne, N. G. Tierney & J. W. Thomson. "Neural Correlates of Self-Face Recognition: An Effect-Location Meta-Analysis." *Brain Research* 1232 (2008): 173–184.
- Puce, A., T. Allison, M. Asgari, J. C. Gore & G. McCarthy. "Differential Sensitivity of Human Visual Cortex to Faces, Letterstrings, and Textures: A Functional Magnetic Resonance Imaging Study." *The Journal of Neuroscience* 16 (1996): 5205–5215.
- Reber, P. J., E. C. Wong & R. B. Buxton. "Comparing the Brain Areas Supporting Nondeclarative Categorization and Recognition Memory." *Cognitive Brain Research* 14 (2002): 245–257.
- Rombouts, S.A.R.B., P. Scheltens, W. Machielsen, F. Barkhof, F. G. C. Hoogenraad, D. J. Veltman, J. Valk & M. P. Witter. "Parametric fMRI Analysis of Visual Encoding in the Human Medial Temporal Lobe." *Hippocampus* 9 (1999): 637–643.
- Sugiura, M., J. Watanabe, Y. Maeda, Y. Matsue, H. Kukuda & R. Kawashima. "Cortical Mechanisms of Visual Self-Recognition." *NeuroImage* 24 (2005): 143–149.
- Taylor, M. J., M. Arsalidou, D. Morris, S. J. Bayless, J. Evans & E. J. Barbeau. "Neural Correlates of Personally Familiar Faces: Parents, Partner and Own Faces." *Human Brain Mapping* 30 (2009): 2008–2020.
- Turak, B., J. Louvel, P. Buser & M. Lamarche. "Event-Related Potentials Recorded from the Cingulate Gyrus During Attentional Tasks: A Study in Patients with Implanted Electrodes." *Neuropsychologia* 40 (2002): 99–107.
- Uddin, L. Q., J. T. Kaplan, I. Molnar-Szakacs, E. Zaidel & M. Iacoboni. "Self-Face Recognition Activates a Frontoparietal 'Mirror' Network in the Right Hemisphere: An Event-Related fMRI Study." *NeuroImage* 25 (2005): 926–935.